arXiv:1006.3915v2 [math.NT] 22 Jun 2010

# TWO IDENTITIES INVOLVING THE CUBIC PARTITION FUNCTION

XINHUA, XIONG

Abstract. We give a new proof of Chan's identity involving the cubic partition function and we also give a new identity for the cubic partition function which is analogues to the Zuckerman's identity for the ordinary partition function.

## 1. INTRODUCTION

Let $p(n)$ be the number of partitions of $n$, defined by $\sum_{n\geq 0} p(n)q^n := \prod_{n=1}^{\infty}(1-q^n)^{-1}$. In connection with his discovery of certain divisibility properties of $p(n)$ Ramanujan stated the identities:

$$\sum_{n=0}^{\infty} p(5n+4)q^n = 5\frac{(q^5;q^5)_\infty^5}{(q;q)_\infty^6}, \tag{1.1}$$

and

$$\sum_{n=0}^{\infty} p(7n+5)q^n = 7\frac{(q^7;q^7)_\infty^3}{(q;q)_\infty^4} + 49q\frac{(q^7;q^7)_\infty^7}{(q;q)_\infty^8}. \tag{1.2}$$

Here and in the rest of the paper we follow the customary $q$-product notation: we set (for $|q| \leq 1$)

$$(a;q)_\infty = \prod_{n=0}^{\infty}(1-aq^n).$$

Both Hardy and MacMahon considered 1.1 as Ramanujan's "Most Beautiful Identity". Darling [7] proved the first and Mordell [9], Watson [11] and Rademacher-Zuckerman [10] gave proofs for both identities. Recently, H.H. Chan and R.P. Lewis [4] also gave different proofs of both identities. All these proofs used the theory of modular functions. In another paper, Zuckerman [12] obtained the following identity (Zuckerman's identity):

$$\begin{aligned}\sum_{n=0}^{\infty} p(25n+24)q^n = 63\cdot 5^2\frac{(q^5;q^5)_\infty^6}{(q;q)_\infty^7} &+ 52\cdot 5^5 q\frac{(q^5;q^5)_\infty^{12}}{(q;q)_\infty^{13}} + 63\cdot 5^7 q^2\frac{(q^5;q^5)_\infty^{18}}{(q;q)_\infty^{19}}\\ &+ 6\cdot 5^{10}q^3\frac{(q^5;q^5)_\infty^{24}}{(q;q)_\infty^{25}} + 5^{12}q^4\frac{(q^5;q^5)_\infty^{30}}{(q;q)_\infty^{31}}.\end{aligned} \tag{1.3}$$

In two recent papers, H.-C. Chan [2, 3] proved a generalization of 1.1 and 1.2 for a certain kind of partition function $a(n)$ which is defined by

$$\sum_{n=0}^{\infty} a(n)q^n := \frac{1}{(q;q)_\infty (q^2;q^2)_\infty}. \tag{1.4}$$

---

2000 *Mathematics Subject Classification.* Primary: 05A15, 05A30 .

Kim [8] noted that $a(n)$ can be interpreted the number of 2-color partitions of $n$ with colors r and g subject to the restriction that the color b appears only in even parts, so he called $a(n)$ to be the cubic partition function owing to the fact that $a(n)$ is related to Ramanujan's cubic continued fraction. Using some identities for the cubic continued fraction, H.C.-Chan derived the following identity :

**Theorem 1.1** ([2] Theorem 1)**.**

$$\sum_{n=0}^{\infty} a(3n+2)q^n = \frac{3(q^3;q^3)_\infty^3 (q^6;q^6)_\infty^3}{(q;q)_\infty^4 (q^2;q^2)_\infty^4}. \tag{1.5}$$

In this note, we only use 3-dissections of functions $\frac{1}{\Phi(-q)}$ and $\frac{1}{\Psi(q)}$ to give a new elementary proof of 1.5 and we also give the following identity for the cubic partition function $a(n)$ which is similar to Zuckerman's identity 1.3 for the cubic partition function $a(n)$.

**Theorem 1.2.**

$$\begin{aligned}\sum_{n=0}^{\infty} a(9n+8)q^n &= 2\cdot 3^3 \frac{(q^3;q^3)_\infty^{30}}{(q;q)_\infty^{19}(q^2;q^2)_\infty^{7}(q^6;q^6)_\infty^{6}} + 8\cdot 3^3 q \frac{(q^3;q^3)_\infty^{21}(q^6;q^6)_\infty^{3}}{(q;q)_\infty^{16}(q^2;q^2)_\infty^{10}} \\ &+ 19\cdot 3^4 q^2 \frac{(q^3;q^3)_\infty^{12}(q^6;q^6)_\infty^{12}}{(q;q)_\infty^{13}(q^2;q^2)_\infty^{13}} - 64\cdot 3^3 q^3 \frac{(q^3;q^3)_\infty^{3}(q^6;q^6)_\infty^{21}}{(q;q)_\infty^{10}(q^2;q^2)_\infty^{16}} \\ &+ 128\cdot 3^3 q^4 \frac{(q^6;q^6)_\infty^{30}}{(q;q)_\infty^{7}(q^2;q^2)_\infty^{19}(q^3;q^3)_\infty^{6}}. \end{aligned} \tag{1.6}$$

Hence $a(9n+8) \equiv 0 \pmod{27}$, which coincides with the result of Chan, he derived this result with different method.

## 2. PRELIMINARIES

We require a few definitions and lemmas. Let

$$\Phi(-q) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} = \frac{(q;q)_\infty^2}{(q^2;q^2)_\infty}, \quad \Psi(q) = \sum_{n=0}^{\infty} q^{(n^2+n)/2} = \frac{(q^2;q^2)_\infty}{(q;q^2)_\infty},$$

and

$$P(q) = \frac{(q^2;q^6)_\infty (q^4;q^6)_\infty (q^3;q^3)_\infty^2}{(q;q)_\infty}, \quad X(-q) = \frac{(q;q)_\infty (q^6;q^6)_\infty^2}{(q^2;q^2)_\infty (q^3;q^3)_\infty}.$$

We will use the following necessaries in proving our main results. The following lemma are the 3-dissections of functions $\frac{1}{\Phi(-q)}$ and $\frac{1}{\Psi(q)}$.

**Lemma 2.1** ([5], last line in the proof Theorem 1)**.**

$$\frac{1}{\Phi(-q)} = \frac{\Phi(-q^9)}{\Phi(-q^3)^4}(\Phi(-q^9)^2 + 2q\Phi(-q^9)X(-q^3) + 4q^2X(-q^3)^2).$$

**Lemma 2.2** ([6], Lemma 2.2)**.**

$$\frac{1}{\Psi(q)} = \frac{\Psi(q^9)}{\Psi(q^3)^4}(P(q^3)^2 - qP(q^3)\Psi(q^9) + q^2\Psi(q^9)^2).$$

**Lemma 2.3.**

$$\Phi(-q)\Psi(q) = (q;q)_\infty (q^2;q^2)_\infty, \quad X(-q)P(q) = (q^3;q^3)_\infty (q^6;q^6)_\infty.$$

*Proof.* We have

$$\Phi(-q)\Psi(q)=\frac{(q;q)_\infty^2}{(q^2;q^2)_\infty}\cdot\frac{(q^2;q^2)_\infty}{(q;q^2)_\infty}=\frac{(q;q)_\infty^2}{(q;q^2)_\infty}=(q;q)_\infty(q^2;q^2)_\infty. \tag{2.1}$$

and

$$\begin{aligned}
X(-q)P(q) &= \frac{(q;q)_\infty(q^6;q^6)_\infty^2}{(q^2;q^2)_\infty(q^3;q^3)_\infty}\cdot\frac{(q^2;q^6)_\infty(q^4;q^6)_\infty(q^3;q^3)_\infty^2}{(q;q)_\infty}\\
&= \frac{(q^2;q^6)_\infty(q^4;q^6)_\infty(q^3;q^3)_\infty(q^6;q^6)_\infty^2}{(q^2;q^2)_\infty}\\
&= \frac{(q^2;q^6)_\infty(q^4;q^6)_\infty(q^3;q^3)_\infty(q^6;q^6)_\infty^2}{(q^2;q^6)_\infty(q^4;q^6)_\infty(q^6;q^6)_\infty}\\
&= (q^3;q^3)_\infty(q^6;q^6)_\infty. \qquad (2.2)
\end{aligned}$$

□

## 3. AN NEW PROOF OF IDENTITY 1.5

We begin with the proof of identity 1.5.

*Proof.* We note that

$$\frac{1}{(q;q)_\infty(q^2;q^2)_\infty}=\frac{(q^2;q^2)_\infty}{(q;q)_\infty^2}\cdot\frac{(q;q^2)_\infty}{(q^2;q^2)_\infty}=\frac{1}{\Phi(-q)\Psi(q)}$$

by using lemma 2.2. So we have

$$\begin{aligned}
&\sum_{n=0}^{\infty}a(n)q^n=\frac{1}{\Phi(-q)\Psi(q)}\\
=\;&\frac{\Phi(-q^9)}{\Phi(-q^3)^4}(\Phi(-q^9)^2+2q\Phi(-q^9)X(-q^3)+4q^2X(-q^3)^2)\\
&\frac{\Psi(q^9)}{\Psi(q^3)^4}(P(q^3)^2-qP(q^3)\Psi(q^9)+q^2\Psi(q^9)^2)\\
=\;&\frac{\Phi(-q^9)\Psi(q^9)}{\Phi(-q^3)^4\Psi(q^3)^4}(\Phi(-q^9)^2P(q^3)^2+2q^3\Phi(-q^9)\Psi(q^9)^2X(-q^3)\\
-\;&4q^3\Psi(q^9)X(-q^3)^2P(q^3)+2q\Phi(-q^9)X(-q^3)P(q^3)^2-q\Phi(-q^9)^2\Psi(q^9)P(q^3)\\
+\;&4q^4\Psi(q^9)^2X(-q^3)^2+q^2\Phi(-q^9)^2\Psi(q^9)^2+4q^2X(-q^3)^2P(q^3)^2\\
-\;&2q^2\Phi(-q^9)\Psi(q^9)X(-q^3)P(q^3)).
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\sum_{n=0}^{\infty}a(3n+2)q^{3n+2}\\
=\;&\frac{\Phi(-q^9)\Psi(q^9)}{\Phi(-q^3)^4\Psi(q^3)^4}(q^2\Phi(-q^9)^2\Psi(q^9)^2+4q^2X(-q^3)^2P(q^3)^2\\
-\;&2q^2\Phi(-q^9)\Psi(q^9)X(-q^3)P(q^3)).
\end{aligned}$$

Dividing by $q^2$ on both sides and replacing $q^3$ by $q$, we obtain

$$\begin{aligned}
&\sum_{n=0}^{\infty}a(3n+2)q^n\\
=\;&\frac{\Phi(-q^3)\Psi(q^3)}{\Phi(-q)^4\Psi(q)^4}(\Phi(-q^3)^2\Psi(q^3)^2+4X(-q)^2P(q)^2-2\Phi(-q^3)\Psi(q^3)X(-q)P(q))
\end{aligned}$$

$$\begin{aligned}
&= \frac{(q^3;q^3)_\infty^3(q^6;q^6)_\infty^3}{(q;q)_\infty^4(q^2;q^2)_\infty^4} + 4\frac{(q^3;q^3)_\infty(q^6;q^6)_\infty}{(q;q)_\infty^4(q^2;q^2)_\infty^4}(q^3;q^3)_\infty^2(q^6;q^6)_\infty^2 \\
&- 2\frac{(q^3;q^3)_\infty^2(q^6;q^6)_\infty^2}{(q;q)_\infty^4(q^2;q^2)_\infty^4}(q^3;q^3)_\infty(q^6;q^6)_\infty \\
&= 3\frac{(q^3;q^3)_\infty^3(q^6;q^6)_\infty^3}{(q;q)_\infty^4(q^2;q^2)_\infty^4},
\end{aligned}$$

by using Lemma 2.3. □

## 4. PROOF OF THE IDENTITY 1.6

We will use the following lemma in proving the identity 1.6.

**Lemma 4.1.** *Let*

$$L := \Phi(-q^9)^2 + 2q\Phi(-q^9)X(-q^3) + 4q^2X(-q^3)^2$$

*and*

$$M := P(q^3)^2 - q\Psi(q^9)P(q^3) + q^2\Psi(q^9)^2.$$

*Then all terms having the exponents of the form of* $3n+2 (n \geq 0)$ *in powers of* $q$ *in* $L^4M^4$ *are* $A+B+C+D+E$*, Where*

$$\begin{aligned}
A &= 40q^2P(q^3)^8X(-q^3)^2\Phi(-q^9)^6 - 32q^2P(q^3)^7X(-q^3)\Phi(-q^9)^7\Psi(q^9) \\
&+ 10q^2P(q^3)^6\Phi(-q^9)^8\Psi(q^9)^2, \\
B &= 512q^5P(q^3)^8X(-q^3)^5\Phi(-q^9)^3 - 1216q^5P(q^3)^7X(-q^3)^4\Phi(-q^9)^4\Psi(q^9) \\
&+ 1280q^5P(q^3)^6X(-q^3)^3\Phi(-q^9)^5\Psi(q^9)^2 \\
&- 640q^5P(q^3)^5X(-q^3)^2\Phi(-q^9)^6\Psi(q^9)^3 \\
&+ 152q^5P(q^3)^4X(-q^3)^3\Phi(-q^9)^7\Psi(q^9)^4 - 16q^5P(q^3)^3\Phi(-q^9)^8\Psi(q^9)^5, \\
C &= 256q^8P(q^3)^8X(-q^3)^8 - 2048q^8P(q^3)^7X(-q^3)^7\Phi(-q^9)\Psi(q^9) \\
&+ 6400q^8P(q^3)^6X(-q^3)^6\Phi(-q^9)^2\Psi(q^9)^2 \\
&- 8192q^8P(q^3)^5X(-q^3)^5\Phi(-q^9)^3\Psi(q^9)^3 \\
&+ 5776q^8P(q^3)^4X(-q^3)^4\Phi(-q^9)^4\Psi(q^9)^4 \\
&- 2048q^8P(q^3)^3X(-q^3)^3\Phi(-q^9)^5\Psi(q^9)^5 \\
&+ 400q^8P(q^3)^2X(-q^3)^2\Phi(-q^9)^6\Psi(q^9)^6 - 32q^8P(q^3)X(-q^3)\Phi(-q^9)^7\Psi(q^9)^7 \\
&+ q^8\Phi(-q^9)^8\Psi(q^9)^8, \\
D &= -4096q^{11}P(q^3)^5X(-q^3)^8\Psi(q^9)^3 + 9728q^{11}P(q^3)^4X(-q^3)^7\Phi(-q^9)\Psi(q^9)^4 \\
&- 10240q^{11}P(q^3)^3X(-q^3)^6\Phi(-q^9)^2\Psi(q^9)^5 \\
&+ 5120q^{11}P(q^3)^2X(-q^3)^5\Phi(-q^9)^3\Psi(q^9)^6 \\
&- 1216q^{11}P(q^3)X(-q^3)^4\Phi(-q^9)^4\Psi(q^9)^7 + 128q^{11}X(-q^3)^3\Phi(-q^9)^5\Psi(q^9)^8
\end{aligned}$$

*and*

$$\begin{aligned}
E &= 2560q^{14}P(q^3)^2X(-q^3)^8\Psi(q^9)^6 - 2048q^{14}P(q^3)X(-q^3)^7\Phi(-q^9)\Psi(q^9)^7 \\
&+ 640q^{14}X(-q^3)^6\Phi(-q^9)^2\Psi(q^9)^8.
\end{aligned}$$

*Proof.* We directly expand the expression of $L^4M^4$ and then extract the terms having exponents $3n+2$ in powers of $q$, then we can obtain the results above. □

**Lemma 4.2.**

$$\begin{aligned}
A &= 2\cdot 3^2q^2\frac{(q^6;q^6)_\infty^6(q^9;q^9)_\infty^{26}}{(q^3;q^3)_\infty^6(q^{18};q^{18})_\infty^{10}}, \\
B &= 8\cdot 3^2q^5\frac{(q^6;q^6)_\infty^3(q^9;q^9)_\infty^{17}}{(q^3;q^3)_\infty^3(q^{18};q^{18})_\infty},
\end{aligned}$$

$$\begin{aligned}
C &= 19\cdot 3^3 q^8 (q^9;q^9)_\infty^8 (q^{18};q^{18})_\infty^8,\\
D &= -64\cdot 3^2 q^{11} \frac{(q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty^{17}}{(q^6;q^6)_\infty^3 (q^9;q^9)_\infty},\\
E &= 128\cdot 3^2 q^{14} \frac{(q^3;q^3)_\infty^6 (q^{18};q^{18})_\infty^{26}}{(q^6;q^6)_\infty^6 (q^9;q^9)_\infty^{10}}.
\end{aligned} \tag{4.1}$$

*Proof.* By using the definitions of $P(q)$, $X(-q)$, the Lemma 2.1, Lemma 2.3 and the formulas

$$(q^6;q^6)_\infty = (q^6;q^{18})_\infty (q^{12};q^{18})_\infty (q^{18};q^{18})_\infty, \quad (q^9;q^9)_\infty = (q^9;q^{18})_\infty (q^{18};q^{18}),$$

we have

$$\begin{aligned}
A &= 40q^2 \frac{(q^6;q^{18})_\infty^8 (q^{12};q^{18})_\infty^8 (q^9;q^9)_\infty^{16} (q^3;q^3)_\infty^2 (q^{18};q^{18})_\infty^4 (q^9;q^9)_\infty^{12}}{(q^3;q^3)_\infty^8 (q^6;q^6)_\infty^2 (q^9;q^9)_\infty^2 (q^{18};q^{18})_\infty^6}\\
&- 32q^2 \frac{(q^6;q^{18})_\infty^7 (q^{12};q^{18})_\infty^7 (q^9;q^9)_\infty^{14} (q^3;q^3)_\infty (q^{18};q^{18})_\infty^2 (q^9;q^9)_\infty^{14} (q^{18};q^{18})_\infty}{(q^3;q^3)_\infty^7 (q^6;q^6)_\infty (q^9;q^9)_\infty (q^{18};q^{18})_\infty^7 (q^9;q^{18})_\infty}\\
&+ 10q^2 \frac{(q^6;q^{18})_\infty^6 (q^{12};q^{18})_\infty^6 (q^9;q^9)_\infty^{12} (q^9;q^9)_\infty^{16} (q^{18};q^{18})_\infty^2}{(q^3;q^3)_\infty^6 (q^{18};q^{18})_\infty^8 (q^9;q^{18})_\infty^2}\\
&= 18q^2 \frac{(q^6;q^6)_\infty^6 (q^9;q^9)_\infty^{26}}{(q^3;q^3)_\infty^6 (q^{18};q^{18})_\infty^{10}}.
\end{aligned}$$

$$\begin{aligned}
B &= 512q^5 \frac{(q^6;q^{18})_\infty^8 (q^{12};q^{18})_\infty^8 (q^9;q^9)_\infty^{16} (q^3;q^3)_\infty^5 (q^{18};q^{18})_\infty^{10} (q^9;q^9)_\infty^6}{(q^3;q^3)_\infty^8 (q^6;q^6)_\infty^5 (q^9;q^9)_\infty^5 (q^{18};q^{18})_\infty^3}\\
&- 1216q^5 \frac{(q^6;q^{18})_\infty^7 (q^{12};q^{18})_\infty^7 (q^9;q^9)_\infty^{14} (q^3;q^3)_\infty^4 (q^{18};q^{18})_\infty^8 (q^9;q^9)_\infty^8 (q^{18};q^{18})_\infty}{(q^3;q^3)_\infty^7 (q^6;q^6)_\infty^4 (q^9;q^9)_\infty^4 (q^{18};q^{18})_\infty^4 (q^9;q^{18})_\infty}\\
&+ 1280q^5 \frac{(q^6;q^{18})_\infty^6 (q^{12};q^{18})_\infty^6 (q^9;q^9)_\infty^{12} (q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty^6 (q^9;q^9)_\infty^{10} (q^{18};q^{18})_\infty^2}{(q^3;q^3)_\infty^6 (q^6;q^6)_\infty^3 (q^9;q^9)_\infty^3 (q^{18};q^{18})_\infty^5 (q^9;q^{18})_\infty^2}\\
&- 640q^5 \frac{(q^6;q^{18})_\infty^5 (q^{12};q^{18})_\infty^5 (q^9;q^9)_\infty^{10} (q^3;q^3)_\infty^2 (q^{18};q^{18})_\infty^4 (q^9;q^9)_\infty^{12} (q^{18};q^{18})_\infty^3}{(q^3;q^3)_\infty^5 (q^6;q^6)_\infty^2 (q^9;q^9)_\infty^2 (q^{18};q^{18})_\infty^6 (q^9;q^{18})_\infty^3}\\
&+ 152q^5 \frac{(q^6;q^{18})_\infty^4 (q^{12};q^{18})_\infty^4 (q^9;q^9)_\infty^8 (q^3;q^3)_\infty (q^{18};q^{18})_\infty^2 (q^9;q^9)_\infty^{14} (q^{18};q^{18})_\infty^4}{(q^3;q^3)_\infty^4 (q^6;q^6)_\infty (q^9;q^9)_\infty (q^{18};q^{18})_\infty^7 (q^9;q^{18})_\infty^4}\\
&- 16q^5 \frac{(q^6;q^{18})_\infty^3 (q^{12};q^{18})_\infty^3 (q^9;q^9)_\infty^6 (q^9;q^9)_\infty^{16} (q^{18};q^{18})_\infty^5}{(q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty^6 (q^9;q^{18})_\infty^5}\\
&= 8\cdot 3^2 q^5 \frac{(q^6;q^6)_\infty^3 (q^9;q^9)_\infty^{17}}{(q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty}.
\end{aligned}$$

$$\begin{aligned}
C &= 256q^8 \frac{(q^6;q^{18})_\infty^8 (q^{12};q^{18})_\infty^8 (q^9;q^9)_\infty^{16} (q^3;q^3)_\infty^8 (q^{18};q^{18})_\infty^{16}}{(q^3;q^3)_\infty^8 (q^6;q^6)_\infty^8 (q^9;q^9)_\infty^8}\\
&- 2048q^8 \frac{(q^6;q^{18})_\infty^7 (q^{12};q^{18})_\infty^7 (q^9;q^9)_\infty^{14} (q^3;q^3)_\infty^7 (q^{18};q^{18})_\infty^{14} (q^9;q^9)_\infty^2 (q^{18};q^{18})_\infty}{(q^3;q^3)_\infty^7 (q^6;q^6)_\infty^7 (q^9;q^9)_\infty^7 (q^{18};q^{18})_\infty (q^9;q^{18})_\infty}\\
&+ 6400q^8 \frac{(q^6;q^{18})_\infty^6 (q^{12};q^{18})_\infty^6 (q^9;q^9)_\infty^{12} (q^3;q^3)_\infty^6 (q^{18};q^{18})_\infty^{12} (q^9;q^9)_\infty^4 (q^{18};q^{18})_\infty^2}{(q^3;q^3)_\infty^6 (q^6;q^6)_\infty^6 (q^9;q^9)_\infty^6 (q^{18};q^{18})_\infty^2 (q^9;q^{18})_\infty^2}\\
&- 8192q^8 \frac{(q^6;q^{18})_\infty^5 (q^{12};q^{18})_\infty^5 (q^9;q^9)_\infty^{10} (q^3;q^3)_\infty^5 (q^{18};q^{18})_\infty^{10} (q^9;q^9)_\infty^6 (q^{18};q^{18})_\infty^3}{(q^3;q^3)_\infty^5 (q^6;q^6)_\infty^5 (q^9;q^9)_\infty^5 (q^{18};q^{18})_\infty^3 (q^9;q^{18})_\infty^3}\\
&+ 5776q^8 \frac{(q^6;q^{18})_\infty^4 (q^{12};q^{18})_\infty^4 (q^9;q^9)_\infty^8 (q^3;q^3)_\infty^4 (q^{18};q^{18})_\infty^8 (q^9;q^9)_\infty^8 (q^{18};q^{18})_\infty^4}{(q^3;q^3)_\infty^4 (q^6;q^6)_\infty^4 (q^9;q^9)_\infty^4 (q^{18};q^{18})_\infty^4 (q^9;q^{18})_\infty^4}\\
&- 2048q^8 \frac{(q^6;q^{18})_\infty^3 (q^{12};q^{18})_\infty^3 (q^9;q^9)_\infty^6 (q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty^6 (q^9;q^9)_\infty^{10} (q^{18};q^{18})_\infty^5}{(q^3;q^3)_\infty^3 (q^6;q^6)_\infty^3 (q^9;q^9)_\infty^3 (q^{18};q^{18})_\infty^5 (q^9;q^{18})_\infty^5}
\end{aligned}$$

$$\begin{aligned}
&+ && 400q^8\frac{(q^6;q^{18})_\infty^2(q^{12};q^{18})_\infty^2(q^9;q^9)_\infty^4(q^3;q^3)_\infty^2(q^{18};q^{18})_\infty^4(q^9;q^9)_\infty^{12}(q^{18};q^{18})_\infty^6}{(q^3;q^3)_\infty^2(q^6;q^6)_\infty^2(q^9;q^9)_\infty^2(q^{18};q^{18})_\infty^6(q^9;q^{18})_\infty^6}\\
&- && 32q^8\frac{(q^6;q^{18})_\infty(q^{12};q^{18})_\infty(q^9;q^9)_\infty^2(q^3;q^3)_\infty(q^{18};q^{18})_\infty^2(q^9;q^9)_\infty^{14}(q^{18};q^{18})_\infty^7}{(q^3;q^3)_\infty(q^6;q^6)_\infty(q^9;q^9)_\infty(q^{18};q^{18})_\infty^7(q^9;q^{18})_\infty^7}\\
&+ && q^8\frac{(q^9;q^9)_\infty^{16}(q^{18};q^{18})_\infty^8}{(q^{18};q^{18})_\infty^8(q^9;q^{18})_\infty^8}\\
&= && 19\cdot 3^3q^8(q^9;q^9)_\infty^8(q^{18};q^{18})_\infty^8.
\end{aligned}$$

$$\begin{aligned}
D &= && -4096q^{11}\frac{(q^6;q^{18})_\infty^5(q^{12};q^{18})_\infty^5(q^9;q^9)_\infty^{10}(q^3;q^3)_\infty^8(q^{18};q^{18})_\infty^{16}(q^{18};q^{18})_\infty^3}{(q^3;q^3)_\infty^5(q^6;q^6)_\infty^8(q^9;q^9)_\infty^8(q^9;q^{18})_\infty^3}\\
&+ && 9728q^{11}\frac{(q^6;q^{18})_\infty^4(q^{12};q^{18})_\infty^4(q^9;q^9)_\infty^8(q^3;q^3)_\infty^7(q^{18};q^{18})_\infty^{14}(q^9;q^9)_\infty^2(q^{18};q^{18})_\infty^4}{(q^3;q^3)_\infty^4(q^6;q^6)_\infty^7(q^9;q^9)_\infty^7(q^{18};q^{18})_\infty(q^9;q^{18})_\infty^4}\\
&- && 10240q^{11}\frac{(q^6;q^{18})_\infty^3(q^{12};q^{18})_\infty^3(q^9;q^9)_\infty^6(q^3;q^3)_\infty^6(q^{18};q^{18})_\infty^{12}(q^9;q^9)_\infty^4(q^{18};q^{18})_\infty^5}{(q^3;q^3)_\infty^3(q^6;q^6)_\infty^6(q^9;q^9)_\infty^6(q^{18};q^{18})_\infty^2(q^9;q^{18})_\infty^5}\\
&+ && 5120q^{11}\frac{(q^6;q^{18})_\infty^2(q^{12};q^{18})_\infty^2(q^9;q^9)_\infty^4(q^3;q^3)_\infty^5(q^{18};q^{18})_\infty^{10}(q^9;q^9)_\infty^6(q^{18};q^{18})_\infty^6}{(q^3;q^3)_\infty^2(q^6;q^6)_\infty^5(q^9;q^9)_\infty^5(q^{18};q^{18})_\infty^3(q^9;q^{18})_\infty^6}
\end{aligned}$$

$$\begin{aligned}
&- && 1216q^{11}\frac{(q^6;q^{18})_\infty(q^{12};q^{18})_\infty(q^9;q^9)_\infty^2(q^3;q^3)_\infty^4(q^{18};q^{18})_\infty^8(q^9;q^9)_\infty^8(q^{18};q^{18})_\infty^7}{(q^3;q^3)_\infty(q^6;q^6)_\infty^4(q^9;q^9)_\infty^4(q^{18};q^{18})_\infty^4(q^9;q^{18})_\infty^7}\\
&+ && 128q^{11}\frac{(q^3;q^3)_\infty^3(q^{18};q^{18})_\infty^6(q^9;q^9)_\infty^{10}(q^{18};q^{18})_\infty^8}{(q^6;q^6)_\infty^3(q^9;q^9)_\infty^3(q^{18};q^{18})_\infty^5(q^9;q^{18})_\infty^8}\\
&= && -64\cdot 3^2q^{11}\frac{(q^3;q^3)_\infty^3(q^{18};q^{18})_\infty^{17}}{(q^6;q^6)_\infty^3(q^9;q^9)_\infty}
\end{aligned}$$

$$\begin{aligned}
E &= && 2560q^{14}\frac{(q^6;q^{18})_\infty^2(q^{12};q^{18})_\infty^2(q^9;q^9)_\infty^4(q^3;q^3)_\infty^8(q^{18};q^{18})_\infty^{16}(q^{18};q^{18})_\infty^6}{(q^3;q^3)_\infty^2(q^6;q^6)_\infty^8(q^9;q^9)_\infty^8(q^9;q^{18})_\infty^6}\\
&- && 2048q^{14}\frac{(q^6;q^{18})_\infty(q^{12};q^{18})_\infty(q^9;q^9)_\infty^2(q^3;q^3)_\infty^7(q^{18};q^{18})_\infty^{14}(q^9;q^9)_\infty^2(q^{18};q^{18})_\infty^7}{(q^3;q^3)_\infty(q^6;q^6)_\infty^7(q^9;q^9)_\infty^7(q^{18};q^{18})_\infty(q^9;q^{18})_\infty^7}\\
&+ && 640q^{14}\frac{(q^3;q^3)_\infty^6(q^{18};q^{18})_\infty^{12}(q^9;q^9)_\infty^4(q^{18};q^{18})_\infty^8}{(q^6;q^6)_\infty^6(q^9;q^9)_\infty^6(q^{18};q^{18})_\infty^2(q^9;q^{18})_\infty^8}\\
&= && 128\cdot 3^2q^{14}\frac{(q^3;q^3)_\infty^6(q^{18};q^{18})_\infty^{26}}{(q^6;q^6)_\infty^6(q^9;q^9)_\infty^{10}}.
\end{aligned}$$

□

Now we prove the identity 1.6.

*Proof.* By Theorem 1.5, Lemma 4.1 and Lemma 4.2, we have

$$\begin{aligned}
\sum_{n=0}^{\infty}a(9n+8)q^{3n+2} &= \text{all terms having the exponents of the form of } 3n+2\\
&\qquad (n\geq 0) \text{ in powers of } q \text{ in } 3\frac{(q^3;q^3)_\infty^3(q^6;q^6)_\infty^3}{(q;q)_\infty^4(q^2;q^2)_\infty^4}\\
&= \text{all terms having the exponents of the form of } 3n+2\\
&\qquad (n\geq 0) \text{ in powers of } q \text{ in } 3\frac{(q^3;q^3)_\infty^3(q^6;q^6)_\infty^3}{\Phi(-q)^4\Psi(q)^4}
\end{aligned}$$

$$\begin{aligned}
&= \text{all terms having the exponents of the form of } 3n+2 \\
&\quad (n \geq 0) \text{ in powers of } q \text{ in } 3(q^3;q^3)_\infty^3 (q^6;q^6)_\infty^3 \\
&\quad \cdot \frac{\Phi(-q^9)^4 \Psi(q^9)^4}{\Phi(-q^3)^{16} \Psi(q^3)^{16}} \cdot L^4 M^4 \\
&= 3(q^3;q^3)_\infty^3 (q^6;q^6)_\infty^3 \cdot \frac{\Phi(-q^9)^4 \Psi(q^9)^4}{\Phi(-q^3)^{16} \Psi(q^3)^{16}} \\
&\quad \cdot (A+B+C+D+E) \\
&= 3\frac{(q^9;q^9)_\infty^4 (q^{18};q^{18})_\infty^4}{(q^3;q^3)_\infty^{16} (q^6;q^6)_\infty^{16}} \cdot (A+B+C+D+E) \\
&= 3\frac{(q^9;q^9)_\infty^4 (q^{18};q^{18})_\infty^4}{(q^3;q^3)_\infty^{16} (q^6;q^6)_\infty^{16}} \cdot \Bigl(18q^2 \frac{(q^6;q^6)^6 (q^9;q^9)^{26}}{(q^3;q^3)^6 (q^{18};q^{18})^{10}} \\
&+ 8 \cdot 3^2 q^5 \frac{(q^6;q^6)_\infty^3 (q^9;q^9)_\infty^{17}}{(q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty} + 19 \cdot 3^3 q^8 (q^9;q^9)_\infty^8 (q^{18};q^{18})_\infty^8 \\
&- 64 \cdot 3^2 q^{11} \frac{(q^3;q^3)_\infty^3 (q^{18};q^{18})_\infty^{17}}{(q^6;q^6)_\infty^3 (q^9;q^9)_\infty} \\
&+ 128 \cdot 3^2 q^{14} \frac{(q^3;q^3)_\infty^6 (q^{18};q^{18})_\infty^{26}}{(q^6;q^6)_\infty^6 (q^9;q^9)_\infty^{10}} \Bigr)
\end{aligned}$$

If we divide $q^2$ on both sides and replace $q^3$ by $q$, we obtain

$$\begin{aligned}
\sum_{n=0}^{\infty} a(9n+8)q^n &= 2 \cdot 3^3 \frac{(q^3;q^3)_\infty^{30}}{(q;q)_\infty^{19} (q^2;q^2)_\infty^7 (q^6;q^6)_\infty^6} + 8 \cdot 3^3 q \frac{(q^3;q^3)_\infty^{21} (q^6;q^6)_\infty^3}{(q;q)_\infty^{16} (q^2;q^2)_\infty^{10}} \\
&+ 19 \cdot 3^4 q^2 \frac{(q^3;q^3)_\infty^{12} (q^6;q^6)_\infty^{12}}{(q;q)_\infty^{13} (q^2;q^2)_\infty^{13}} - 64 \cdot 3^3 q^3 \frac{(q^3;q^3)_\infty^3 (q^6;q^6)_\infty^{21}}{(q;q)_\infty^{10} (q^2;q^2)_\infty^{16}} \\
&+ 128 \cdot 3^3 q^4 \frac{(q^6;q^6)_\infty^{30}}{(q;q)_\infty^7 (q^2;q^2)_\infty^{19} (q^3;q^3)_\infty^6},
\end{aligned}$$

which is the identity 1.6. □

## 5. ACKNOWLEDGEMENT

The author thanks Professor Liu Zhi-Guo's helpful comments.

## References

1. B.C. Berndt, *Ramanujan's Notebooks, Part III*, Springer-Verlag, New York, 1991.
2. H.-C. Chan, *Ramanujan's cubic continued fraction and a generalization of his "most beautiful identity"*, Int. J. Number Theory, **6** (2010), 673-680
3. H.-C. Chan, *A new proof of two identities involving Ramanujan's cubic continued fraction*, Ramanujan J. **21** (2010), 173–180.
4. H.H. Chan and R.P. Lewis, *Partition identities and congruences associated with the Fourier coefficients of the Euler products*, J. Computational and Applied Mathematics, **160** (2003), 69-75
5. M.D. Hirschhorn and J.A. Sellers, *Arithmetic relations for overpartitions*, Journal of Combinatorial Mathematics and Combinatorial Computing, **53** (2005), 65-73.
6. M.D. Hirschhorn and J.A. Sellers, *Arithmetic properties of partitions with odd parts distinct*, Ramanujan J. online first, 2010.
7. Darling, *Proofs of certain identities and congruences enunciated by S. Ramanujan*, Proc. London Math. Soc. (2), **19** (1921), 350-372.

8. B. Kim, *A crank analog on a certain kind of partition function arising from the cubic continued fraction*, preprint, 2008.
9. Mordell, *Note on certain modular relations considered by Messrs. Ramanujan, Darling, and Rogers*, Proc. London Math. Soc. (2), **20** (1922), 408-416.
10. H. Radermacher and H.S. Zuckerman, *A new proof of Ramanujan's identities*, Ann. of Math., **40** (1939), 473-489.
11. G.N. Watson, *Beweis von Ramanujans Vermutungen über Zerfällungsanzahlen*, J. Reine und Angew. Math., **179** (1938), 97-128.
12. H.S. Zukerman, *Identities analogous to Ramanujan'a identities involving the partition function*, Duke Math.J. **5** (1939), 88-110.

Department of Mathematics, China Three Gorges University, Yichang, Hubei Province, 443002, P.R. China

*E-mail address*: `xinhuaxiong@ctgu.edu.cn`